



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Making the equations homogeneous by writing in them $z = 1$,

$$[(x^2 + y^2) - (a^2 + b^2)z^2] \tan^2 \varphi = 4z^2(b^2x^2 + a^2y - a^2b^2z^2). \quad (C)$$

$$\frac{zx}{a^2} + \frac{\lambda y}{b^2} = z. \quad (D)$$

Eliminating z , the coordinates of the points in which (C) and (D) intersect are given by the equation

$$\left[(x^2 + y^2) - (a^2 + b^2) \left(\frac{zx}{a^2} + \frac{\lambda y}{b^2} \right)^2 \right]^2 \tan^2 \varphi = 4 \left(\frac{zx}{a^2} + \frac{\lambda y}{b^2} \right)^2 \times \left[b^2x^2 + a^2y - a^2b^2 \left(\frac{zx}{a^2} + \frac{\lambda y}{b^2} \right)^2 \right].$$

Expanding and reducing, we get

$$Ax^4 + Bx^3y + Cx^2y^2 + Dxy^3 + Ey^4 = 0. \quad (E)$$

Where

$$A = b^4[a^2 - b^2(a^2 + b^2)x^2] \tan^2 \varphi - 4a^2b^8x^2 + 4a^2b^{10}x^4,$$

$$B = 4[a^2b^6(a^2 + b^2)^2x^3\lambda - a^4b^4(a^2 + b^2)x\lambda] \tan^2 \varphi - 8a^4b^6x\lambda + 16a^4b^8x^3\lambda,$$

$$C = [2a^4b^4 - 2a^2b^2(a^2 + b^2)(a^4\lambda^2 + b^4x^2) + 6a^4b^4(a^2 + b^2)^2x^2\lambda^2] \tan^2 \varphi - 4(a^6b^4\lambda^2 + a^4b^6x^2) + 24a^6b^6x^2\lambda^2,$$

$$D = 4[a^6b^2(a^2 + b^2)^2x\lambda^3 - a^4b^4(a^2 + b^2)x\lambda] \tan^2 \varphi - 8a^6b^4 + 16a^8b^4x\lambda^2,$$

$$E = a^4[b^2 - a^2(a^2 + b^2)\lambda^2] \tan^2 \varphi - 4a^8b^2\lambda^2 + 4a^{10}b^2\lambda^4.$$

But if (D) is tang't to (C), equation (E) must have equal roots and therefore its discriminant vanishes. Therefore

$$4(12AE - 3BD + C^2)^3 = (72ACE + 9BCD - 27AD^2 - 27EB^2 - 2C^3)^2.$$

(Salmons Higher Algebra, 3rd edition, p. 306.)

This is an equation between the coordinates x , λ and is, therefore the required envelope.

NOTE ON AN INDETERMINATE EQN., BY WM. HOOVER, A. M.—The following quotation is from a communication by M. E. Catalan to *Journal de Mathematiques* for August, 1882.

“The identity

$$(a + b)^2 (a - 2b)^2 (b - 2a)^2 + 27a^2b^2 = 4(a^2 - ab + b^2)^3,$$

easy of verification, gives an indefinite number of solutions, in entire numbers, of

$$x^2 + 3y^2 = z^3.$$

“We can take

$$x = \frac{1}{2}(a+b)(a-2b)(b-2a), y = \frac{3}{2}ab(a-b), \text{ and } z = a^2 - ab + b^2.”$$